

Is GARCH(1,1) as good a model as the Nobel prize accolades would imply?¹

Cătălin Stărică ²

First draft: 13 October 2003

This version: 2 November 2003

¹This research has been supported by The Bank of Sweden Tercentenary Foundation.

²Department of Mathematical Statistics, Chalmers University of Technology, S-412 96, Gothenburg, Sweden. Email: starica@math.chalmers.se. Web page: www.math.chalmers.se/~starica

Abstract

This paper investigates the relevance of the stationary, conditional, parametric ARCH modeling paradigm as embodied by the GARCH(1,1) process to describing and forecasting the dynamics of returns of the Standard & Poors 500 (S&P 500) stock market index.

A detailed analysis of the series of S&P 500 returns featured in Section 3.2 of the Advanced Information note on the Bank of Sweden Prize in Economic Sciences in Memory of Alfred Nobel reveals that during the period under discussion, there were no (statistically significant) differences between GARCH(1,1) modeling and a simple non-stationary, non-parametric regression approach to next-day volatility forecasting. A second finding is that the GARCH(1,1) model severely over-estimated the unconditional variance of returns during the period under study.

For example, the annualized implied GARCH(1,1) unconditional standard deviation of the sample is 35% while the sample standard deviation estimate is a mere 19%. Over-estimation of the unconditional variance leads to poor volatility forecasts during the period under discussion with the MSE of GARCH(1,1) 1-year ahead volatility more than 4 times bigger than the MSE of a forecast based on historical volatility.

We test and reject the hypothesis that a GARCH(1,1) process is the true data generating process of the longer sample of returns of the S&P 500 stock market index between March 4, 1957 and October 9, 2003. We investigate then the alternative use of the GARCH(1,1) process as a local, stationary approximation of the data and find that the GARCH(1,1)

model fails during significantly long periods to provide a good local description to the time series of returns on the S&P 500 and Dow Jones Industrial Average indexes.

Since the estimated coefficients of the GARCH model change significantly through time, it is not clear how the GARCH(1,1) model can be used for volatility forecasting over longer horizons. A comparison between the GARCH(1,1) volatility forecasts and a simple approach based on historical volatility questions the relevance of the GARCH(1,1) dynamics for longer horizon volatility forecasting for both the S&P 500 and Dow Jones Industrial Average indexes.

JEL classification: C14, C16, C32.

Keywords and Phrases: stock returns, volatility, Garch(1,1), non-stationarities, unconditional time-varying volatility, IGARCH effect, longer-horizon forecasts.

1. INTRODUCTION

Even a casual look through the econometric literature of the last two decades reveals a drastic change in the conceptual treatment of economic time series. The modeling of such time series moved from a static set-up to one that recognizes the importance of fitting the time-varying features of macro-economic and financial data.

In particular, it is now widely accepted that the covariance structure of returns (referred often as volatility) changes through time. A large part of the modern econometric literature frames modeling of time-varying volatility in the *autoregressive conditional heteroskedastic* (ARCH) framework, a stationary, parametric, conditional approach that postulates that the main time-varying feature of returns is the *conditional covariance structure*³ while assuming in the same time that *the unconditional covariance remains constant through time* (see for example, the survey Bollerslev et al [4]). While the autoregressive conditional heteroskedastic approach to modeling time-varying volatility is currently prevalent, alternative methodologies for volatility modeling exist in the econometric literature. In particular, the non-stationary framework that assumes the *unconditional variance* to be the main time-varying feature of returns has a long tradition that can be traced back to Officer [15], Hsu, Miller and Wichern [11], Merton [12], French et al. [7].

This paper is motivated by the desire to better understand the relevance of the stationary, parametric conditional ARCH paradigm *as embodied by the GARCH(1,1) process* to modeling and forecasting the returns of financial indexes. We focus on the GARCH(1,1)

³The conditional covariance structure is supposed to follow an autoregressive mechanism from where the name of the paradigm.

process (Bollerslev [3], Taylor [17]) since this model is widely used and highly regarded in practice as well as in the academic discourse. Producing GARCH(1,1) time-varying volatility estimates is part of the daily routine in many financial institutions. In the academic literature, the GARCH(1,1) process seems to be perceived as a *realistic data generating process for financial returns*. As a result, a large number of econometric and statistical papers that develop estimation and testing techniques⁴ based on the assumption of a GARCH-type data generating mechanism, use the GARCH(1,1) model to actually implement their results.

The main goal of the paper is to investigate how close is the simple endogenous dynamics imposed by a GARCH(1,1) process to the true dynamics of returns of main financial indexes. To this end we analyze in detail the log-returns of the S&P 500 stock market index between March 4, 1957 and October 9, 2003⁵. Further evidence is brought in from the analysis of the Dow Jones Industrial Average index covering the same period.

Our endeavor is, by nature, limited in scope. An analysis of the level of detail characterizing ours cannot encompass, in the length of one paper, alternative ARCH-type models and/or more series. Hence, the conclusions we draw are fortuitously limited to one model

⁴This includes testing of economic and financial theories like the Arbitrage Pricing Theory or the Capital Asset Pricing Model.

⁵The S&P 500 stock index took the present form of an average of the price of 500 stocks on March 4, 1957.

and two time series. However it is worth remembering that the model is the most celebrated member of the ARCH family while the time series represent the epitome of financial return data.

The paper is organized as follows. Section 2 focuses on modeling and forecasting issues related to the series of S&P 500 returns between May 16, 1995 to April 29, 2003. In Section 2.1 the stationary, parametric GARCH(1,1) model is compared to a simple non-stationary, non-parametric regression approach in the context of next-day volatility forecasting. Section 2.2 discusses the GARCH(1,1) estimated unconditional variance of returns during the period under study. The impact of these estimates on GARCH(1,1) forecasting is analyzed in Section 2.3.

Section 3 discusses modeling and forecasting issues related to the longer series of S&P returns between March 4, 1957 and October 9, 2003. In section 3.1 we test the hypothesis that a GARCH(1,1) process is the true data generating process of the longer sample of returns on the S&P 500 and Dow Jones Industrial Average stock market indexes. In Section 3.2 we offer a possible explanation for the poor estimates of the unconditional volatility documented in Section 2.2. We also evaluate the GARCH(1,1) process as a local stationary approximation of the return data. In Section 3.3 a comparison between the GARCH(1,1) volatility forecasts and a simple approach based on historical volatility evaluates the relevance of the GARCH(1,1) dynamics for longer horizon volatility forecasting for both the S&P 500 and Dow Jones Industrial Average indexes. Section 4 concludes.

2. THE GARCH(1,1) MODEL AND THE S&P 500 INDEX RETURNS BETWEEN MAY 16, 1995 AND APRIL 29, 2003

When modeling the returns on the S&P 500 index in the stationary, parametric, conditional ARCH framework, the *working assumption* is often that the data generating process is the GARCH(1,1) model

$$(2.1) \quad r_t = z_t h_t^{1/2}, \quad h_t = \alpha_0 + \alpha_1 r_{t-1}^2 + \beta_1 h_{t-1},$$

where (z_t) are iid, $Ez = 0$, $Ez^2 = 1$. Since the GARCH(1,1) process is stationary, assuming it as data generating process implicitly assumes that return data is stationary.

We begin with a detailed analysis of the series of S&P 500 returns between May 16, 1995 to April 29, 2003 - 2000 observations in all⁶. The upper panel of Figure 2.1 shows the daily logarithmic returns (first differences of the logarithms of daily closing price) (on the bottom row the longer sample of returns on the S&P 500 index from March 4, 1957 to October 9, 2003 is displayed⁷). According to the Advanced Information note, fitting a conditionally normal GARCH(1,1) process to the series on the first row in Figure 2.1 yields the estimated parameters $\hat{\alpha}_0 = 2 \times 10^{-6}$, $\hat{\alpha}_1 = 0.091$, and $\hat{\beta}_1 = 0.899$ corresponding to a

⁶This particular time series is featured in the illustration of the use of the GARCH(1,1) model in estimating and forecasting volatility in Section 3.2 of the Advanced Information note on the Bank of Sweden Prize in Economic Sciences in Memory of Alfred Nobel. The note is available as

<http://www.nobel.se/economics/laureates/2003/eoadv.pdf>

⁷Since most of the analyses in the sequel are done under the assumption of stationarity, the returns of the week starting with 19 October 1987 were considered exceptional and were removed from the sample.

value of the likelihood of 6136.2. Our estimates were slightly different

$$(2.2) \quad \hat{\alpha}_0 = 1.4264 \times 10^{-6}, \quad \hat{\alpha}_1 = 0.0897, \quad \hat{\beta}_1 = 0.9061.$$

and correspond to a higher value of the likelihood of 6136.6⁸. Note that $\hat{\alpha}_1 + \hat{\beta}_1 = 0.995$, a value very close to 1.




Figure 2.1. Top: S&P 500 returns between May 16, 1995 and April 29, 2003 (top) and from March 4, 1957 to October 9, 2003 (bottom).

⁸For the rest of the discussion we stick with our estimates although using the estimates from Advanced Information note does not change in any way the results.

2.1. Next-day volatility forecasting. One of the main achievements of the GARCH modeling consists in providing accurate next-day volatility forecasts. Statistically, this statement is supported by measures of the goodness of fit of the model (2.1) based on the estimated innovations or residuals defined as:

$$(2.3) \quad \hat{z}_t = r_t / \hat{h}_t^{1/2}, \quad \hat{h}_t = \hat{\alpha}_0 + \hat{\alpha}_1 r_{t-1}^2 + \hat{\beta}_1 \hat{h}_{t-1}, \quad t = 1, \dots, n.$$

Residuals that are close to being independent are taken as evidence of accurate next-day volatility forecasts. Figure 2.2 displays the estimated GARCH(1,1) volatility as well as the residuals from the model (2.1) with parameters (2.2) corresponding to the period from May 16, 1995 to April 29, 2003.

While the sample ACF of the absolute returns displays significant linear dependency at lags as large as 100 (Figure 2.3–top), the absolute residuals pass the Ljung-Box test of independence for at least the first 100 lags (Figure 2.3–bottom)(for more details on the Ljung-Box test, see Brockwell and Davis [5]).




Figure 2.2. Top: Annualized volatility S&P 500 from May 16, 1995 to April 29. Bottom: Estimated innovations.

For the data under analysis, the GARCH(1,1) volatility admits the following infinite moving average approximation:

$$\begin{aligned}
 \hat{h}_t &= \hat{\alpha}_0 + \hat{\alpha}_1 r_{t-1}^2 + \hat{\beta}_1 \hat{h}_{t-1} \\
 &= \hat{\alpha}_0 + \hat{\alpha}_1 r_{t-1}^2 + \hat{\beta}_1 (\hat{\alpha}_0 + \hat{\alpha}_1 r_{t-2}^2 + \hat{\beta}_1 \hat{h}_{t-2}) \\
 &= \hat{\alpha}_0 (1 + \hat{\beta}_1) + \hat{\alpha}_1 (r_{t-1}^2 + \hat{\beta}_1 r_{t-2}^2) + \hat{\beta}_1^2 \hat{h}_{t-2} \\
 &\quad \dots \\
 &= \hat{\alpha}_0 / (1 - \hat{\beta}_1) + \hat{\alpha}_1 (r_{t-1}^2 + \hat{\beta}_1 r_{t-2}^2 + \hat{\beta}_1^2 r_{t-3}^2 + \dots) \\
 &\approx \hat{\alpha}_0 / (1 - \hat{\beta}_1) + (1 - \hat{\beta}_1) (r_{t-1}^2 + \hat{\beta}_1 r_{t-2}^2 + \dots) \\
 (2.4) \quad &\approx \frac{r_{t-1}^2 + \hat{\beta}_1 r_{t-2}^2 + \hat{\beta}_1^2 r_{t-3}^2 + \dots}{1 + \hat{\beta}_1 + \hat{\beta}_1^2 + \dots}.
 \end{aligned}$$




Figure 2.3. Sample ACF of absolute values of returns (top) and of absolute values of residuals (bottom-left). The p -values of the Ljung-Box statistic for the absolute values of the residuals (bottom-right). The hypothesis of independence is rejected for small p -values.

The first approximation consists in replacing $\hat{\alpha}_1$ with $1 - \hat{\beta}_1$ (recall that $\hat{\alpha}_1 + \hat{\beta}_1 = 0.995$) while the second one neglects the term $\hat{\alpha}_0 / (1 - \hat{\beta}_1)^9$. Figure 2.4 displays the conditional variance process as well as its approximation given by (2.4) and show a close correspondence between the two processes.

⁹While more important during the low volatility period up to the beginning of 1997, this term accounts on average, for 10% of the variance during the period 1997-2003.




Figure 2.4. GARCH(1,1) conditional variance (full line) and the approximation in (2.4) (dotted line). The broken line marks the contribution of the term $\hat{\alpha}_0/(1 - \hat{\beta}_1)$. The paths of the two processes are, most of the time, very close to each other.

The approximation (2.4) is, simply, an exponential smoother, the exponentially weighted moving average (EWMA) applied to (r_t^2) . It is known that EWMA forecasting is optimal (in the mean-square error sense)¹⁰ for the state space model

$$(2.5) \quad x_t = \mu_t + \epsilon_t,$$

$$(2.6) \quad \mu_t = \mu_{t-1} + \nu_t,$$

¹⁰The optimality follows from the fact that EWMA reduces to the Kalman filter in this case.

where (ϵ_t) and (ν_t) are iid sequences, $E\epsilon = 0$, $E\nu = 0$. In the state space model set-up, equation (2.6) incorporates the uncertainty about the form of the model.

Note that the GARCH(1,1) model can be written in the form of equation (2.5):

$$r_t^2 = h_t + h_t(z_t^2 - 1) := h_t + \epsilon_t,$$

with $E\epsilon = 0$. Going one step further,¹¹ equation (2.6) would translate to

$$(2.7) \quad h_t = h_{t-1} + \nu_t.$$

In words, equation (2.7) acknowledges that the dynamic of the variance is unpredictable. While a small variance of the noise ν can imply that the daily changes are so small as to be ignored, over longer periods of time the movements of the variance cannot be foreseen hence modeled.

A closely related set-up, subtly different but which shares with the state model representation the ability to incorporate our uncertainty about the form of the model is that of the non-parametric regression (see for example, Wand and Jones [18]). The mentioned uncertainty is handled in this case by modeling the signal μ as a deterministic function of time¹²

$$(2.8) \quad x_t = \mu(t) + \epsilon_t,$$

¹¹The discussion here is purely at a formal level. No modeling will be done in the space state set-up. Instead a closely related framework to address the uncertainty of the model will be used. Had we pursued the state space specification, special care would have been needed to assure that h_t stays positive

¹²In both situations the local level or the local trend can be estimated as well as desired.

where (ϵ_t) is an iid sequence, $E\epsilon = 0$.

This association suggests the following simple non-parametric, non-stationary alternative approach. Assume that

$$(2.9) \quad r_t = \sigma(t) \epsilon_t, \quad E\epsilon = 0, \quad E\epsilon^2 = 1.$$

where $\sigma(t)$ is a deterministic, smooth function of time. In words, assume that the *unconditional* variance is time-varying and that the returns are independent (but of course, not identically distributed). Rewrite equation (2.9) in the form of (2.8)

$$(2.10) \quad r_t^2 = \sigma^2(t) + \sigma^2(t)(\epsilon_t^2 - 1) := \sigma^2(t) + \epsilon_t, \quad E\epsilon = 0.$$

Following the non-parametric regression literature (see for example, Wand and Jones [18]), an estimate of the time-varying unconditional variance $\sigma^2(t)$ is given by

$$(2.11) \quad \hat{\sigma}^2(t; b) = \sum_{k=1}^n W_k(t; b) r_k^2$$

where b is the *bandwidth* and the weights W_k are given by

$$(2.12) \quad W_k(t; b) = K\left(\frac{k-t}{b}\right) / \sum_{k=1}^n K\left(\frac{k-t}{b}\right),$$

where K is a kernel (for details on the definition and the properties of a kernel function see the reference above). The expressions (2.11), (2.12) are the Nadaraya-Watson, or zero-degree local polynomial, kernel estimate.

The specific of the problem at hand is that we want to *forecast* the next-day volatility, i.e. use the return data available at time $t - 1$ to forecast $\sigma(t)$. This is accomplished by the right choice of a kernel in (2.12). More concretely, the kernel K will be a *half-kernel* or

a *one-sided kernel*, i.e. $K(x) = 0$ for $x \geq 0$. This implies that $W_k(t; b) = 0$ if $k \geq t$. With this choice of a kernel, equation (2.11) yields the forecasted time-varying variance

$$(2.13) \quad \hat{\sigma}^2(t; b) = \sum_{k=1}^{t-1} K\left(\frac{k-t}{b}\right) r_k^2 / \sum_{k=1}^{t-1} K\left(\frac{k-t}{b}\right).$$

In the sequel we use the exponential and the normal half-kernels as illustrations. Note that $\hat{\sigma}^2(t; b)$, the estimated time-varying unconditional variance at time t depends only on information prior to this moment, i.e. only the returns up to time $t - 1$ have been used in estimation.




Figure 2.5. The average of the squared residuals (ASR) defined in (2.14) as a function of b for the two choices of kernel: the exponential (on the left) and the normal (on the right). A minimum is attained at $b_{ASR}^{exp} = 11$ and $b_{ASR}^{normal} = 12$, respectively.

The key quantity in the non-parametric regression is b , the bandwidth. From the plethora of methods proposed in the extensive statistical literature on the bandwidth selection (see for example Gasser et al. [8]), the simplest one takes b to minimize the average of the

squared residuals (ASR) of the previous one-step-ahead forecast

$$(2.14) \quad \hat{b}_{ASR} = \operatorname{argmin}_{b \geq 0} ASR(b), \quad ASR(b) := n^{-1} \sum_{i=1}^n (\hat{\sigma}^2(t; b) - r_t^2)^2.$$




Figure 2.6. Sample ACF of the absolute values of residuals from non-parametric estimation of volatility using the exponential (top) and normal (bottom) half-kernels. The absolute values display no linear dependence and the hypothesis of independence is not rejected based on a Ljung-Box test at any lag smaller than 100.

In the non-parametric regression literature this approach corresponds to the well-studied *cross-validation* method for the particular case of a half-kernel. Its asymptotic properties can be obtained by extending the results of Chu and Marron ([6]) to half-kernels (see also

Gijbels et al. [9]). Figure 2.5 displays the ASR as a function of b for two choices of kernel: the exponential (on the left) and the normal (on the right). Both graphs have a convex shape attaining a minimum at $b_{ASR}^{exp} = 11$ and $b_{ASR}^{normal} = 12$, respectively¹³.

Figure 2.6 displays the sample ACF of the residuals from the non-parametric regression

$$(2.15) \quad \hat{\varepsilon}_t = r_t / \hat{\sigma}(t; \hat{b}_{ASR}), \quad t = 1, \dots, n.$$

The residuals show no linear dependence and the hypothesis of independence is not rejected based on a Ljung-Box test at any lag smaller than 100.

Figure 2.7 displays the GARCH(1,1) volatility together with the non-parametric estimate (2.13) with a normal half-kernel. The paths of the two volatility processes follow each other closely.




Figure 2.7. (Annualized) Volatility estimates: GARCH(1,1) version (full line) and the non-parametric regression estimate (2.13) with a normal half-kernel (dotted line). The paths of the two volatility processes are, most of the time, very close.

¹³In the notation of equation (2.4), the value $b_{ASR}^{exp} = 11$ corresponds to a $\beta = 0.91$.

The evidence presented shows that, during the period under discussion, there were no (statistically significant) differences between GARCH(1,1) and a simple non-stationary, non-parametric regression approach to next-day volatility forecasting.

An objection against the non-parametric framework outlined above could be that it lacks any dynamics and would be embodied by the question: What has this approach to say about the future?¹⁴. Furthermore, one could argue that, by contrast, the GARCH(1,1) model, by specifying a certain endogenous mechanism for the volatility process, is capable to foresee future developments in the movements of prices. Shortly, the GARCH(1,1) model is preferable because it has a vision of the future.

In the sequel we will analyze the relevance of this possible objection/argument from a number of different perspectives. First we will discuss some modeling and forecasting implications of postulating a GARCH(1,1) volatility mechanism on the returns between May 16, 1995 to April 29, 2003 (sections 2.2 and 2.3). Then we will take a closer look at the relevance of the GARCH(1,1) dynamics for the longer S&P 500 sample covering the period between March 4, 1957 and October 9, 2003 (sections 3.1, 3.2). Finally, both the

¹⁴The answer to this question, that might not fully satisfy the ones in favor of tight parametric modeling, is: since we assume that the volatility is driven by exogenous factors (that are hard to identify and hence we leave unspecified), i.e. we do not know if it will move up or down, and that it evolves smoothly and slowly, i.e. it will take a while until significant changes will be noticed, the best we can do is to accurately measure the current level and to forecast it as the return volatility for the close future. The near future returns will be modeled as iid with a variance equal to today's estimate.

GARCH(1,1) methodology and the non-parametric regression approach will be considered in the frame of longer horizon volatility forecasts in section 3.3.

2.2. Modeling performance. As mentioned earlier, a common working assumption in the financial econometric literature is that the GARCH(1,1) process (2.1) is the true data generating process for financial returns. Under such an assumption and given a data sample, a GARCH(1,1) process with parameters estimated on a reasonably long subsample¹⁵ of the data should provide a good model for the whole sample. More concretely, the GARCH(1,1) process with parameters (2.2) estimated on the 2000 observation long subsample¹⁶ displayed on the top of Figure 2.1 should provide a reasonable description also for the 11727 observation long sample on the bottom of the same figure¹⁷. To check if this is the case, we display in Figure 5.1 24 randomly generated samples of length 11727 from the model (2.1) with parameters (2.2) together with the plot of the true return series. Note that the over-all aspect of the simulated samples is quite different from that of the real data.

Figure 5.2 displays the squared values of the samples in Figure 5.1. Differences between the aspect of the simulated samples and that of the return series are strongly visible. In particular, the variance of the real data appears to be smaller.

¹⁵The subsample size should guarantee that the estimation error is likely to be small.

¹⁶A sample size of 2000 is commonly assumed to be sufficient for a precise estimation of a GARCH(1,1) model.

¹⁷11727 is the the length of the sample from March 4, 1957 to October 9, 2003.

The visual impression from Figure 5.1 and 5.2 is confirmed by simulations. The support of the simulated¹⁸ distribution of the sample variance for samples of length 11727 from the model (2.2) was the interval [0.00014,0.01526]. The value of the variance of the returns in Figure 2.1 (bottom graph) is 0.00008.

This simple simulation exercise highlights the need for a closer look at the GARCH(1,1) estimated unconditional variance. The model (2.1) was first estimated on the subsample from May 16, 1995 to April 30, 1999 (the first 1000 observations in the sample) and then re-estimated every 5 days on a sample that contains all past observations, i.e. all observations between May 16, 1995 and the date of the re-estimation. Under the assumption of weak stationarity, i.e. $\alpha_1 + \beta_1 < 1$, the unconditional variance of the GARCH(1,1) model (2.1) is given by

$$(2.16) \quad \sigma^2 := \alpha_0 / (1 - \alpha_1 - \beta_1).$$

Figure 2.8 displays the annualized estimated unconditional GARCH(1,1) standard deviation (sd) together with the annualized sample sd, i.e. the square root of 250 times the average of square returns from May 16, 1995 to current time location. The graph shows a sd from 1.5 to 5 times bigger than the sample sd¹⁹. Hence, during the period under discussion, the unconditional variance point estimates of a GARCH(1,1) model are severely out-of-line with the sample point estimates of the unconditional variance of returns.

¹⁸25,000 samples were simulated.

¹⁹For example, the parameter values in the Advanced Information note imply an annualized sd of 35% while the sample annualized sd is of merely 19%.




Figure 2.8. Estimated GARCH (1,1) sd (2.16) (dotted line) together with sample sd (both estimates are annualized) (full line). The model (2.1) is re-estimated every 5 days on a sample that contains all past observations beginning from May 16, 1995. The time mark corresponds to the end of the sub-sample that yields the two sd estimates. The graph shows a big discrepancy between the two estimates.

Since for the model (2.1) the volatility forecast at longer horizons is, practically, the unconditional variance (see equation (2.17)), failing to produce accurate point estimates for this last quantity will, most likely, produce poor longer horizon volatility forecasts. We will now investigate this issue.

2.3. Forecasting performance. Let us now evaluate the out-of-sample forecasting performance of the GARCH(1,1) model based on the sample on top of Figure 2.1.

Assuming a GARCH(1,1) data generating process (2.1) that also satisfy²⁰ $\alpha_1 + \beta_1 < 1$, it follows that the minimum Mean Square Error forecast for Er_{t+p}^2 , the return variance h -steps ahead, is

$$(2.17) \quad \sigma_{t+p}^{2, GARCH} := E_t r_{t+h}^2 = \sigma^2 + (\alpha_1 + \beta_1)^p (h_t - \sigma^2),$$

where σ^2 is the unconditional variance defined in (2.16). Note that, since $\alpha_1 + \beta_1 < 1$, for large h , the forecast $\sigma_{t+p}^{2, GARCH}$ is the unconditional variance, σ^2 .

The minimum MSE forecast for $E(r_{t+1} + \dots + r_{t+p})^2$, the variance of the next p aggregated returns, is then given by

$$(2.18) \quad \bar{\sigma}_{t,p}^{2, GARCH} := E_t (r_{t+1} + \dots + r_{t+p})^2 = \sigma_{t+1}^{2, GARCH} + \dots + \sigma_{t+p}^{2, GARCH}.$$

We take as benchmark (BM) for volatility forecasting the simple non-stationary model (2.9). Since no dynamics is specified for the variance, future observations r_{t+1}, r_{t+2}, \dots are modeled as iid with constant variance $\hat{\sigma}^2(t)$, an estimate of $\sigma^2(t)$. In the sequel we use the sample variance of the previous year of returns as the estimate for $\sigma^2(t)$. The forecast for Er_{t+p}^2 is then given by

$$(2.19) \quad \sigma_{t+p}^{2, BM} := \widehat{Er}_t^2 = \hat{\sigma}^2(t),$$

²⁰If this condition is not fulfilled, the GARCH(1,1) process, if stationary, has infinite variance.




Figure 2.9. The (future) realized sd (annualized) at horizon 250 (full line) together with GARCH forecast (broken line) and the benchmark (dotted line). The time mark is the beginning of the period over which the forecast is made.

The forecast for $E(r_{t+1} + \dots + r_{t+p})^2$, the variance of the next h aggregated returns is simply

$$(2.20) \quad \bar{\sigma}_{t,p}^{2, BM} := p \hat{\sigma}^2(t) = \frac{p}{250} \sum_{i=1}^{250} r_{t-i+1}^2.$$

The GARCH(1,1) model is estimated initially on the first 1000 data points corresponding to the interval from May 16, 1995 to April 30, 1999, and re-estimated every 5 days (every week). Contemporaneously, $\sigma^2(t)$ is estimated as the average of past 250 square returns. After every re-estimation, volatility forecasts are made for the year to come ($p = 1, \dots, 250$) using (2.18) and (2.20). The data used in the out-of-sample variance forecasting comparison covers the interval May 3, 1999 to April 29, 2003.




Figure 2.10. The ratio MSE^{GARCH}/MSE^{BM} . The MSE^* are defined in (2.22). On the x -axis, h , the forecast horizon.

Define the following measure of the realized volatility in the interval $[t+1, t+p]$

$$(2.21) \quad \bar{r}_{t,p}^2 := \sum_{i=1}^h r_{t+i}^2,$$

We calculated and compared the following MSE

$$(2.22) \quad MSE^*(p) := \sum_{t=1}^n (\bar{r}_{t,p}^2 - \bar{\sigma}_{t,p}^{2,*})^2$$

with $*$ standing for BM or GARCH. The MSE (2.22) is preferred to the simpler MSE

$$\sum_{t=1}^n (r_{t+p}^2 - \sigma_{t+p}^{2,*})^2$$

since this last one uses a poor measure of the realized return volatility²¹. Through averaging some of the idiosyncratic noise in the daily squared return data is canceled yielding (2.21), a better measure against which to check the quality of the two forecasts.

Figure 2.9 shows the extent of the impact on forecasting of the over-estimation of unconditional variance illustrated in Figure 2.8. Note also the similarity of the shape of the GARCH curves in Figure 2.8 and Figure 2.9.

Both Figure 2.10 and 2.9 show that a simple model with a *time-varying unconditional variance* (hence non-stationary) produced better out-of-sample forecasting than the GARCH(1,1) model in the period May 3, 1999 to April 29, 2003.

In this section, we have seen that the GARCH(1,1) process produced point estimates of the unconditional variance that were severely out-of-line with the sample point estimates during the period May 3, 1999 to April 29, 2003. As a consequence, during this period the model yielded poor longer horizon forecasts. The analysis we just concluded raises a number of questions. Does the GARCH(1,1) process always over-estimate unconditional variances? Or was the period we analyzed a special one? And if yes, in what way? To answer these questions we performed a detailed analysis of the fit of the GARCH(1,1) on the time series of returns of the S&P 500 index between March 4, 1957 and October 9, 2003. The results are presented in the next section.

²¹It is well known (see Andersen and Bollerslev [1]) that the realized square returns are poor estimates of the day-by-day movements in volatility, as the idiosyncratic component of daily returns is large.

3. THE GARCH(1,1) MODEL AND THE S&P 500 INDEX RETURNS BETWEEN MARCH 4, 1957 AND OCTOBER 9, 2003

We begin with a re-examination of the working assumptions of the previous analysis. The fact that the GARCH(1,1) process is a *stationary* model²² raises up front a methodological choice since one uses the GARCH(1,1) model differently, depending on the working assumptions one is willing to make about the data to be modeled. Were the econometrician convinced that the data is stationary, she would assume that a GARCH(1,1) process is the true data generating process and she would estimate a GARCH(1,1) model on the whole data set. She would try always to use as much data as possible in order to support any statistical statement or to make forecasts. Had she reasons to believe that the data is non-stationary, then the stationary GARCH(1,1) process could possibly become a useful local approximation of the true data generating process. The parameters of the model should then be re-estimated periodically on a moving window. Also she would shy away from making statistical statements based on long samples and in forecasting she would prefer to use the most recent past for the calibration of the model. However, were the data unconditional features time-varying, it is not clear of what help would the GARCH(1,1) model then be in producing volatility forecasts over longer horizons.

As we see, the two working assumptions on the nature of the data at hand have very different methodological implications both on the estimation of the GARCH(1,1) model as

²²The stationarity is central to issues of statistical estimation and of forecasting.

well as on its use in forecasting. In the sequel we investigate in detail the behavior of the GARCH(1,1) model in the two outlined frameworks.

3.1. Working hypothesis: The returns are stationary. The GARCH(1,1) is the true data generating process. The working assumption in this section will be that the returns on the S&P 500 index between March 4, 1957 and October 9, 2003 form a stationary time series. In this set-up one can test the hypothesis that a given parametric model is the data generating process of this time series. More concretely, the estimated parameters of such a model should be statistically the same no matter what portion of the data is used in estimation. In particular, the parameters should be statistically identical were they estimated on an increasing sample or on a window that moves through the data. Detection of significant changes in the values of the estimated parameters leads to rejecting the parameter model as the data generating process.

A GARCH(1,1) process was fit to the S&P 500 series of returns between March 4, 1957 and October 9, 2003 yielding the following parameters

$$(3.1) \quad \hat{\alpha}_0 = 4.78 \times 10^{-7}, \quad \hat{\alpha}_1 = 0.0696, \quad \hat{\beta}_1 = 0.9267.$$

To evaluate the suitability of a GARCH(1,1) model as data generating process for the S&P 500 return series, the model (2.1) was initially estimated on the first 2000 observations of the sample in Figure 2.1 (bottom) corresponding roughly to the period 1957-1964. Then the model was re-estimated every 50 observations both on a sample containing 2000 past observations and on a sample containing all past observations. The results of the estimation




Figure 3.1. Estimated fourth moment of the residuals \hat{z} for S&P 500 data. The model was initially estimated on the first 2000 observations of the sample corresponding roughly to the period 1957-1964, then re-estimated every 50 observations on a sample containing 2000 past observations (left) and all past observations (right).

are displayed in Figure 5.3. The confidence intervals correspond to the quasi-maximum likelihood (QML) estimation method and take into account possible misspecifications of the conditional distribution (Gouriéroux [10], Berkes et al. [2], Straumann and Mikosch [16]). The graphs in Figure 5.3 show pronounced instability of the estimated values of the parameters.

The asymptotic normality of the QML estimators depends on the existence of the finite fourth moment for the distribution of the innovations z (assumption N3 of Theorem 7.1 Straumann and Mikosch [16]). Figure 3.1 displays the fourth moment of the estimated residuals for the S&P 500 data together with the 95% confidence intervals (under the assumption of independent innovations) and seems to confirm that the fourth moment of the innovations is finite.

A similar picture emerges from analyzing the returns on the Dow Jones Industrial stock index. The evolution of the estimated GARCH(1,1) coefficients displayed in Figure 5.4 follow a similar pattern to the one in Figure 5.3.




Figure 3.2. The histogram of the number of estimated overlapping confidence intervals for the parameters α_1 and β_1 in a group of 5 independent samples of size 2000 from a GARCH (1,1) model with parameters (3.1). 25000 groups were simulated. On the left, the common histogram, on the right the probabilities on a log-scale. A value of -4 in this graph corresponds to an event that will happen with a probability of 10^{-4} . The two sets of histograms, i.e. one corresponding to α_1 and the other to β_1 are, practically, identical.

The hypothesis that a GARCH(1,1) process is the data generating process can be formally tested as follows. Five subsamples of length 2000 separated one from the next by 400 observations²³ were used to produce 95% confidence intervals. Our test statistic is

²³The first subsample consist of the first 2000 data points. 400 observations are dropped and the second subsample starts with observation 2401 and contains the next 2000 observations, etc. The length

the number of pairs of overlapping confidence intervals. Under the null of a common GARCH(1,1) data generating process, the test statistic should be close to 10. A small number of overlapping pairs constitute evidence against the null. For parameter α_1 , 2 out of the 10 possible pairs of confidence intervals do overlap, while for parameter β_1 the number of overlapping pairs is 4. The probability of this two events, i.e. the probability of observing less than m overlapping confidence intervals out of 10 possible pairs, $m = 2$ and $m = 4$, can be bounded as follows. The probability that two $100 \times \alpha\%$ confidence intervals overlap is greater than the probability that both the intervals contain the true parameter, i.e. α^2 . Assuming that the actual coverage of the confidence intervals in Figure 5.3 is the theoretical one²⁴, i.e. 95%, the probability of seeing less than or 2 overlapping intervals out of 10 possible pairs can be easily calculated as being *at most* 3×10^{-7} while the probability of the separating blocks was chosen such that it maximizes the distance between the subsamples, yielding nevertheless 5 non-overlapping series. 2 consecutive samples were separated by more than one year and a half of data to guarantee that the assumption of independent subsamples is likely to be reasonable.

²⁴The test based on the confidence intervals of parameter α_1 still rejects the null hypothesis at 5% if the actual level of coverage is in fact as low as 71% and at 1% if the actual level is of 78% or higher. The test based on the confidence intervals of parameter β_1 rejects the hypothesis of stationarity at 5% if the actual level of coverage is in fact as low as 84% and at 1% if the actual level is of 88% or higher. Note that this calculations yield rather conservative, i.e. high, probabilities for the event of interest, which is observing less than m overlapping confidence intervals out of 10 possible pairs, as they are based on a rough bound on the probability that the confidence intervals of two independent samples overlap and not on the actual value of this probability. The simulation study shows that for a coverage of around 91%, the probability of the event of interest is in fact smaller than 4×10^{-5} for $m = 2$, and smaller than 10^{-4} for $m = 4$, respectively.

of seeing less than or 4 overlapping intervals out 10 possible pairs is *at most* 1.4×10^{-4} . To further investigate the issue of the actual coverage level, a simulation study was performed where 12000 samples of size 2000 from a GARCH(1,1) process with parameters estimated on the long series of S&P 500 returns (3.1) were generated. The innovations were drawn from the empirical distribution of the residuals of the model (2.1) with parameters given by (3.1). A GARCH(1,1) model was estimated on every simulated sample and the 95% confidence intervals for the 3 parameters were constructed. The actual coverage of the 95% confidence intervals so constructed was of 94.9% for α_0 , 90.9% for α_1 and 91.8% for β_1 . Furthermore, 25000 groups of five independent samples of size 2000 each were formed and the number of overlapping confidence intervals in a group counted. Figure 3.2 displays the results. Since the actual coverage levels for the parameters α_1 and β_1 are so close, the two sets of histograms, i.e. one corresponding to α_1 and the other to β_1 , are practically identical and hence we show only the one corresponding to α_1 . As a conclusion, even when taking into account the possibly lower level of coverage of the estimated confidence intervals, the null is strongly rejected.

For the Dow Jones Industrial index the hypothesis that parameter α_1 is constant through time is rejected with a p -value of at most 0.0014 while the hypothesis of constant β_1 was rejected with a p -value of at most 0.0101.

To get a visual feeling for the type of results one gets in the case when a GARCH(1,1) model is really the data generating process, let us have a look at Figure 5.6 which displays the outcome of the estimation of GARCH(1,1) parameters for a simulated sample from a

GARCH(1,1) model. The sample that is 11727 observations long, is displayed in Figure 5.5 together with the real S&P 500 data. The parameters of the data generating process were (3.1). The estimation was done on a 2000 observation long window that moves through the data and hence the graphs correspond to the ones on the left of Figure 5.3. Figure 5.6 reveals a very different behavior of the estimated parameters without the clear time-evolution present in Figure 5.3. Note that the true parameters are always inside the confidence intervals.




Figure 3.3. Estimated GARCH (1,1) standard deviation (2.16) (dotted line) together with sample standard deviation (both estimates are annualized) (full line). The model (2.1) is re-estimated every 50 days on a sample including all past observations. The time mark corresponds to the end of the sub-sample that yields the two standard deviation estimates. The two curves wander always far apart.

We end this section with a comparison between the point estimates of the GARCH(1,1) unconditional variance and of the sample variance. Under the assumption of stationary returns, the values of the parameters estimated on a growing sample containing all past returns were used to produce the GARCH(1,1) unconditional sd (the broken line in Figure 3.3). The corresponding sample sd is depicted by the full line in the same figure. The two curves wander wide apart suggesting that the GARCH(1,1) process estimated on an increasing sample produces poor point estimates of the variance.

The evidence presented in this section indicates that *the GARCH(1,1) process is not the data generating process for the returns of the S&P 500 stock index between March 4, 1957 and October 9, 2003.*

The significant changes of the value of the estimated parameters in Figure 5.3 suggests that a more realistic working hypothesis is that of non-stationary data (in particular, the unconditional variance could be time-varying). If this was the case, the GARCH(1,1) model might turn out to, at least, provide good local stationary approximation to the process of index returns. This issue is investigated in the next section.

3.2. Working hypothesis: The returns are non-stationary. The GARCH(1,1) process is a local stationary approximation of the changing data. Working under the assumption that unconditional features of the data might be also time-varying allows us to provide a possible explanation for the poor unconditional variance point estimates produced by the GARCH(1,1) model during the period May 16, 1995 to April 29, 2003 documented in section 2.1. The analysis in this section will also individuate all other

periods when the GARCH(1,1) model yielded point estimates of the unconditional variance that were not in line with the sample unconditional variance and hence, failed to provide a good local stationary approximation of the true data generating process of returns²⁵.

Figure 5.7 displays the estimated $\alpha_1 + \beta_1$ under the assumption of non-stationary data. The model (2.1) has been initially estimated on the first 2000 observations of the sample in Figure 2.1 (bottom) corresponding roughly to the period 1957-1964, then re-estimated every 50 observations on a sample containing 2000 past observations. The graph shows that the IGARCH effect significantly²⁶ affects the GARCH(1,1) models (estimated on a sample that ends) during the period 1997-2003²⁷. This fact at its turn, is likely to cause the explosion of the estimated unconditional variance of the GARCH(1,1) processes fitted on samples that end during this period (see (2.18)).

To see that indeed this is the case, let us take look at the bottom graph of the same Figure 5.7 where the GARCH(1,1) unconditional sd (broken line) and the corresponding sample sd (full line) are displayed. The GARCH(1,1) unconditional sd is obtained using (2.16) from the values of the parameters estimated on a window of size 2000 moving through the data and that are displayed in Figure 5.3. The graph shows a good agreement between

²⁵A good local approximation model should fit reasonably well relatively short subsamples. In the sequel we individuate only the periods when the GARCH(1,1) model does not provide an accurate description of the unconditional variance. Of course, there could be that other stochastic features of the data are not well-captured during the periods when the estimation of the unconditional variance is satisfactory.

²⁶The point estimate is close to 1 and, more importantly, 1 belongs to the 95% one-sided confidence interval.

²⁷During the interval 1994-1996, the value 1 is the upper bound of the confidence interval.

the two estimates at all times except during the period when the IGARCH effect becomes strongly statistically significant, i.e. samples that end in the interval 1997-2003^{28,29}.

The bottom graph in Figure 5.7 together with the analysis in Section 2 show that the GARCH(1,1) model fails to provide a local stationary approximation to the time series of returns on the S&P 500 during significantly long periods.

An analysis of the Dow Jones Industrial Average stock index produces similar findings. The results are displayed in Figures 5.8.

An explanation for the strong IGARCH effect in the second half of the 90's can be the sharp change in the unconditional variance (see Mikosch and Starica [13]). There it is proved, both theoretically and empirically, that sharp changes in the unconditional variance can cause the IGARCH effect. Figure 3.4 displays non-parametric estimates of the unconditional sd together with the 95% confidence intervals³⁰ for the S&P 500 returns (top) and the Dow Jones industrial index returns (bottom). The two graphs show a pronounced

²⁸The analysis was also performed with smaller sample sizes of 1500, 1250 and 1000. As expected, the confidence intervals in Figures 5.7 and 5.8 get wider and hence less meaningful. However, for every sample sized mentioned, there is always a period between 1997 and 2003 where the unconditional variance of the estimated model explodes. Estimation based on samples smaller than 1000 observations is infeasible as it produces extremely unstable coefficients and renders problematic the use of any asymptotic result.

²⁹Contrast this finding with the statement on page 16 of the Advanced Information note: "Condition $\alpha_1 + \beta_1 < 1$ is necessary and sufficient for the first-order GARCH process to be weakly stationary, and the estimated model (on the short S&P 500, n.n.) satisfies this condition."

³⁰The method used to obtain the estimates is that of kernel smoothing in the framework of non-parametric regression with non-random equi-distant design points. For more details on the performance of this method on financial data see Mikosch and Starica [14].

increase of the volatility from around 5% in 1993-1994 to three times as much (around 15%) in the period 2000-2003.




Figure 3.4. Estimated unconditional standard deviation (annualized) with 95% confidence intervals for the S&P 500 returns (top) and Dow Jones returns (bottom). The shaded areas correspond to bear market periods.

3.3. The relevance of the GARCH(1,1) model for longer horizon forecasting of volatility of index return series. As we have seen in section 3.1, the coefficients of the

GARCH(1,1) model change significantly through time. This phenomena raises naturally the question about the relevance of the model for volatility forecasting over longer horizons. This issue is investigated in the sequel by comparing the forecasting performance of a GARCH(1,1) with that of a simple approach based on historical volatility. Both the S&P 500 data set and the Down Jones Industrial stock index returns are analyzed.




Figure 3.5. The ratio MSE^{GARCH}/MSE^{BM} . The MSE^* are defined in (2.22). The GARCH(1,1) model has been re-estimated on an increasing sample containing all past observations (left) and on a moving window of 2000 past observations (right). On the x -axis, p , the forecast horizon.

The set-up is the one described in Section 2.3. A GARCH(1,1) model was estimated on the first 2000 observations corresponding roughly to the period 1957-1964. Then the model was re-estimated every 10 observations (every two business weeks) both on a sample containing 2000 past observations and on a sample containing all past observations. Every time the model was re-estimated, GARCH volatility forecast (see equation (2.18)) as well

as forecasts based on historical volatility (see equation (2.20)) were made for horizons from 1 day to 250 days ahead. The MSE for the two volatility forecasts were calculated using (2.22) and the ratio MSE^{GARCH}/MSE^{BM} is displayed in Figure 3.5. While for shorter (than 60 days ahead for locally estimated model and than 40 days ahead for a model estimated on all past data) horizons, the GARCH(1,1) volatility forecast has a smaller MSE, for longer forecasting horizons the historical volatility does significantly better.




Figure 3.6. The ratio MSE^{GARCH}/MSE^{BM} . The MSE^* are defined in (2.22). The GARCH(1,1) model has been re-estimated on an increasing sample containing all past observations (left) and on a moving window of 2000 past observations (right). On the x -axis, p , the forecast horizon.

The results displayed in the graphs in Figure 3.5 seem to question the relevance of the GARCH(1,1) dynamics for longer horizon volatility forecast of the S&P 500 return series.

The finding that the GARCH(1,1) dynamics provides a poor longer horizon forecasting basis (when compared to a simple non-stationary approach centered on historical volatility)

documented for the S&P 500 return time series in Figure 3.5 was confirmed by similar results for the Dow Jones Industrial stock index returns (see Figure 3.6).




Figure 3.7. The realized 1-year ahead volatility (full line) together with the GARCH(1,1) forecasted volatility (dotted line) (top) for the S&P 500 data. The broken line in the bottom graph is the historical volatility forecast based on the past 250 returns (all quantities are annualized).

Finally, Figure 3.7 displays the realized 1-year ahead volatility, the GARCH(1,1) forecasted volatility and the historical volatility forecast based on the past 250 returns (all the quantities are annualized). The graphs show a poor coincidence between the realized and

the forecasted GARCH(1,1) volatilities. Notably, the GARCH(1,1) forecast becomes especially volatile (even when compared with the forecast based on historical volatility) during the period when the estimation of the model is significantly affected by the IGARCH effect, i.e. the period that starts towards the end of 1997 and lasts to the end of the sample.

4. CONCLUSIONS

In this paper we investigated how truthful is the simple endogenous volatility dynamics imposed by a GARCH(1,1) process to the evolution of returns of main financial indexes. Our analysis concentrated first, on the series of S&P 500 returns between May 16, 1995 to April 29, 2003³¹ and was extended then to the longer series of S&P returns between March 4, 1957 and October 9, 2003.

The analysis of the shorter sample showed no (statistically significant) differences between a GARCH(1,1) modeling and a simple non-stationary, non-parametric regression approach to next-day volatility forecasting.

We tested and rejected the hypothesis that a GARCH(1,1) process is the data generating process for the series of returns on the S&P 500 stock index from March 4, 1957 to October 9, 2003. QML estimation of the parameters of a GARCH(1,1) model on a window that moves through the data produced statistically different coefficients. Since the parameters of the GARCH(1,1) model change significantly through time, it is not clear how the model can be used for volatility forecasting over longer horizons.

³¹This series is featured in Section 3.2 of the Advanced Information note on the Bank of Sweden Prize in Economic Sciences in Memory of Alfred Nobel.

We also evaluated the behavior of the GARCH(1,1) model as a local, stationary approximation for the data. We have found that the IGARCH effect is not innocuous as it is often claimed. In fact, it seems that during the periods when the IGARCH effect is statistically significant, the forecasting performance of the GARCH(1,1) model deteriorates drastically at all horizons. Analyzing one of these periods, we found the MSE for longer horizon GARCH(1,1) volatility forecasts to be up to more than four times bigger than a simple forecast based on historical volatility.

In a forecasting comparison with a simple non-stationary approach centered on historical volatility, the finding that the GARCH(1,1) dynamics provides a poor longer horizon forecasting basis was documented for the S&P 500 return time series as well as for the Dow Jones Industrial Average index returns.

Acknowledgement: The constructive criticism of Holger Drees, Thomas Mikosch and Richard Davis has benefited the paper, both in its conception and in its presentation.

5. APPENDIX: BIG FIGURES




Figure 5.1. 24 randomly generated samples of length 11727 from the model (2.1) with parameters (2.2) (first 24 graphs). The S&P 500 returns from March 4, 1957 to October 9, 2003 (bottom-right). The aspect of the real data is different from that of the simulated samples.

REFERENCES

- [1] ANDERSEN, T, BOLLERSLEV, T. (1998) Answering the critics: Yes, ARCH models do provide good volatility forecasts. *International Economic Review* **39**, 885–905.
- [2] BERKES, I., HORVATH, L. AND KOKOSZKA, P. (2002) Probabilistic and statistical properties of GARCH processes. To appear *Fields Institute Communications*. Available at <http://www.math.usu.edu/~piotr>.
- [3] BOLLERSLEV, T. (1986) Generalized autoregressive conditional heteroskedasticity. *J. Econometrics* **31**, 307–327.
- [4] BOLLERSLEV, T., ENGLE, R.F. AND NELSON, D.B. (1994) GARCH models. In: Engle, R.F. and McFadden, D.L. (Eds.) *Handbook of Econometrics*. Vol. **4**, pp. 2961–3038. Elsevier, Amsterdam.
- [5] BROCKWELL AND DAVIS, R. (1987) *Time Series: Theory and Methods*. Second Edition. Springer, New York.
- [6] CHU, C. AND MARRON, J. (1991) Comparison of two bandwidth selectors with dependent errors. *Ann. Statist.* **19**, 1906–1918.
- [7] FRENCH, K., SCHWERT, W. AND STAMBAUGH, R. (1987) Expected stock returns and volatility. *J. Financial Economics* **19**, 3–29.
- [8] GASSER, T., ENGEL, J. AND SEIFERT, B. (1993) Nonparametric function estimation. In: *Computational Statistics. Handbook of Statistics*. Vol. **9**, pp. 423–465. North-Holland, Amsterdam.
- [9] GIJBELS, I., POPE, A., AND WAND, M. P. (1999) Understanding exponential smoothing via kernel regression. *J. R. Statist. Soc. B*, **51**, 39–50.
- [10] GOURIÉROUX, C. (1997) *ARCH Models and Financial Applications*. Springer, New York.
- [11] HSU, D.A., MILLER, R., AND WICHERN, D. (1974) On the stable Paretian behavior of stock-market prices. *J. of American Statistical Association* **69**, 108–113.
- [12] MERTON, R. (1980) On estimating the expected return on the market: an exploratory investigation. *J. Financial Economics* **8**, 323–361.
- [13] MIKOSCH, T. AND STĂRICĂ, C. (2004) Non-stationarities in financial time series, the long-range dependence and IGARCH effects. *Rev. Econom. Statist.* To appear. Available under www.math.ku.dk/~mikosch.
- [14] MIKOSCH, T. AND STĂRICĂ, C. (2003) Stock market risk-return inference. An unconditional non-parametric approach. Technical Report. Available under www.math.ku.dk/~mikosch.
- [15] OFFICER, R. (1976) The variability of the market factor of the New York Stock Exchange. *J. Business* **46**, 434–453.
- [16] STRAUMANN, D. AND MIKOSCH, T. (2003) Quasi-MLE in heteroscedastic times series: a stochastic recurrence equations approach. Technical Report. Available under www.math.ku.dk/~mikosch.
- [17] TAYLOR, S.J. (1986) *Modelling Financial Time Series*. Wiley, Chichester.
- [18] WAND, M.P. AND JONES, M.C. (1995) *Kernel Smoothing*. Chapman and Hall, London.




Figure 5.2. The squares of 24 randomly generated samples of length 11727 from the model (2.1) with parameters (2.2) (first 24 graphs). The squares of the S&P 500 returns from March 4, 1957 to October 9, 2003. The aspect of the real data is different from that of the simulated samples.




Figure 5.3. Estimated coefficients of the model (2.1) for S&P 500 returns. The model was initially estimated on the first 2000 observations of the sample in Figure 2.1 (bottom) corresponding roughly to the period 1957-1964, then re-estimated every 50 observations on a sample containing 2000 past observations (left column) and on a sample containing all the past observations. The right column display the results of both estimations. The confidence intervals take into account possible misspecifications of the conditional distribution (Gourièroux [10], Berkes et al. [2], Straumann and Mikosch [16]). The dotted lines in the graphs on the left are the parameters estimated on the whole sample. The graphs show pronounced instability of the estimated values of the parameters.




Figure 5.4. Estimated coefficients of the model (2.1) for Dow Jones Industrial Average returns. The model was initially estimated on the first 2000 observations of the sample in Figure 2.1 (bottom) corresponding roughly to the period 1957-1964, then re-estimated every 50 observations on a sample containing 2000 past observations (left column) and on a sample containing all the past observations. The right column displays the results of both estimations. The confidence intervals take into account possible misspecifications of the conditional distribution (Gouriéroux [10], Berkes et al. [2], Straumann and Mikosch [16]). The dotted lines in the graphs on the left are the parameters estimated on the whole sample. The graphs show pronounced instability of the estimated values of the parameters.




Figure 5.5. Top: Simulated S&P 500 returns (top). The sample has been obtained from GARCH(1,1) process with parameters (3.1) estimated on the sample from March 4, 1957 to October 9, 2003 (bottom).




Figure 5.6. Estimated coefficients of the model (2.1) on the simulated data in Figure 5.5. The model has been initially estimated on the first 2000 observations of the simulated sample corresponding roughly to the period 1957-1964, then re-estimated every 50 observations on a sample containing 2000 past observations. The confidence intervals take into account possible misspecifications of the conditional distribution (Gourièroux [10], Berkes et al. [2], Straumann and Mikosch [16]). The true parameters are always inside the confidence bands.




Figure 5.7. Top: Estimated $\alpha_1 + \beta_1$ with one-sided 95% confidence interval. The confidence interval takes into account possible misspecifications of the conditional distribution. Bottom: Estimated GARCH (1,1) sd (2.16) (dotted line) together with sample sd (both estimates are annualized) (full line) for the S&P 500 log-returns. The set-up is that of Figure 5.3. The time mark corresponds to the end of the sub-sample that yields the two standard deviation estimates. While most of the time the two curves in the bottom graph are remarkably close to each other, the GARCH(1,1) variance seems to explode towards the end of the sample.




Figure 5.8. Top: Estimated $\alpha_1 + \beta_1$ with one-sided 95% confidence interval. The confidence interval takes into account possible misspecifications of the conditional distribution.. Bottom: Estimated GARCH (1,1) sd (2.16) (dotted line) together with sample sd (both estimates are annualized) (full line) for the Dow Jones Industrial Average log-returns. The set-up is that of Figure 5.4. The time mark corresponds to the end of the sub-sample that yields the two standard deviation estimates. While most of the time the two curves in the bottom graph are remarkably close to each other, the GARCH(1,1) variance seems to explode towards the end of the sample.